



Mr. Andrew  
Chamblor

Rotator Cuff Repair  
Rehabilitation  
Guidelines  
**(Arthroscopic/mini-  
open/open procedures)**

# ROTATOR CUFF REPAIR

## (Arthroscopic/mini-open/open procedures)

### INTRODUCTION

---

The guidelines that follow are a frame work of basic exercises and management strategies based on the patient who has had a **LARGE** cuff tear (>5-6cm) repaired. The size of the rotator cuff tear and the strength of the repair of the tendon back to the bone vary from patient to patient. Patients who have had a small tear repaired **may** be progressed faster as tolerated.

#### **CLASSIFICATION OF ROTATOR CUFF TEARS:**

SMALL = Less than 1cm; MEDIUM = 1-3cms; LARGE = 3-5cms;

MASSIVE = greater than 5cms

The physiotherapy programme will need to be **individualised** for each patient, all exercises should be performed without pain and the details of specific restrictions will be in the post-operative instructions. If you have not received these please ring the consultant's secretary.

Emphasise to the patient the importance of protecting the repair to allow soft-tissue healing in the first two phases. Alongside this we aim to regain passive range of movement (PROM) and good scapulohumeral rhythm (SHR) before moving onto strengthening in stage 3. The milestones may be used to assess whether you feel the patient is making good progress or not.

Shoulder rehabilitation is more than strength-training of the shoulder muscles alone. The (neuromuscular) rehabilitation addresses the whole shoulder girdle, upper extremity, core stability and training of the kinetic chain.

### NOTES

---

**Subscapularis Repair:** if the subscapularis muscle is repaired, the range of movement into lateral rotation needs to be limited to neutral and no resisted internal rotation should be taught during the first 6-8 weeks to protect the repair (check the post op note).

**Abduction pillow/brace:** larger tears have a greater risk of re-tear due to their size, often poor tissue quality and the repair being under greater tension. In these patients, the emphasis is significantly more on protecting the repair in the early stages rather than regaining range of movement; therefore they **may** be managed in an abduction brace initially.

**Open Rotator Cuff Repairs:** the open procedure requires release of the deltoid muscle from the acromion. The anterior-mid fibres of deltoid therefore need to be protected from full active use **until 6-8 weeks** post operatively.

**InSpace™ implantation (Ballon):** please see accompanying OrthoSpace InSpace™ Implantation protocol and <https://www.youtube.com/watch?v=cejMRmNnmJc> for more information

### POST SURGERY

---

## Phase I (1 – 14 days)

### Goals:

- Maintain integrity of the repair
  - Sling at all times except while dressing/washing or doing exercises
  - Teach sling, dressing and personal hygiene techniques
- Management of pain, inflammation and muscle inhibition
  - Analgesics, NSAID's, ice, sling, passive movement and posture
- Teach shoulder girdle control/setting and relaxation
  - Retraction and depression
- Gradually increase PROM **as tolerated/not into pain/do not force or stretch**
  - Pendulum with good SHR
  - ER/IR (between 0°-45° of scaption with elbow supported)
- Hand, wrist, elbow and neck range of movement (ROM) exercises as required
- Advice on sleeping position
  - Wearing sling, if supine use a pillow beneath the elbow to prevent the shoulder resting in extension
- Align developing collagen fibres and prevent muscle atrophy
  - Sub-maximal, pain-free isometrics in neutral (<30% maximal voluntary contraction) as tolerated

### Precautions:

- Sling usually for 6 weeks (dependent on size of the tear – see post-op instructions)
- No active movement of elbow away from body
- No lifting of objects
- No excessive shoulder extension
- No excessive stretching or sudden movements
- No supporting of body weight on operated upper limb
- Keep wounds clean and dry

### Milestones:

Pain, inflammation and muscle inhibition well managed  
Return to pre-operative sleep patterns  
Good scapula setting  
PROM: ER to neutral and Flexion 45° with good SHR

□

□

- □ **Phase 2 (15 days – 6 weeks)**

### Goals:

- Allow healing of soft tissue – do not over-load healing tissue
- Continue to manage and reduce pain, inflammation and muscle inhibition
  - As phase I
  - Alternate treatment strategies as appropriate e.g. manual therapy techniques, taping
- Once stitches have been removed begin scar massage
- Gradually restore full, pre-op PROM **as tolerated/not into pain/do not force or stretch**
  - Refer to post-op note for specific restrictions

- Introduce active-assisted range of movement (AAROM) from three weeks **as tolerated/not into pain**
  - Refer to post-op note for specific restrictions
- Active range of movement (AROM):
  - In large tears, it will take up to six weeks before it is safe to introduce AROM, progressing from AAROM **as tolerated/not into pain**
  - In small tears, AROM may be possible from 4 weeks progressing from AAROM **as tolerated/not into pain**
  - Refer to post-op note for specific restrictions
- Re-establish dynamic shoulder stability
  - Ensure good SHR through PROM and use it to guide progression of AAROM and AROM

#### Precautions:

- No lifting
- No excessive shoulder extension
- No excessive stretching or sudden movements
- No supporting of body weight by hands
- Usually wean out of the sling after 6 weeks (dependent on size of the tear – see post-op instructions)

#### Milestones at 3 weeks:

- Pain, inflammation and muscle inhibition well managed
- Passive flexion aim to 50% of pre-operative range (dependant on size of tear)
- Passive external rotation to neutral
- Good SHR with PROM

#### Milestones at 6 weeks:

- Pain, inflammation and muscle inhibition well managed
- Passive flexion aim to 100% of pre-operative range (dependant on size of tear)
- Passive external rotation to 30°
- Good SHR with PROM

### Phase 3 (7 - 14 weeks)

#### Goals:

- Full PROM
  - Introduce multi-directional stretching into end of range **as tolerated/not into pain**
- Gradually restore pre-operative AAROM and then AROM with good SHR **as tolerated/not into pain**

- Gradually introduce and progress shoulder strengthening **as tolerated/not into pain**
  - Introduce scapular stabilisation exercises
    - Introduce cuff strengthening, progressing to maximal isometrics then to isotonic strengthening
      - Introduce proprioceptive exercises
    - Gradually progress to shoulder and upper limb strengthening **as long as the patient is able to elevate the arm without the shoulder/scapular 'hitching'**
    - Gradually progress to dynamic and rhythmic stabilisation exercises
- Gradually progress lower limb and core strengthening as required
- Gradually return to light, non-repetitive functional activities
- Gradually introduce light, early-stage sport-specific exercises
- Return to driving (right 6-8 weeks, left 8-10 weeks) smaller tears maybe able to start earlier

#### Precautions:

- Avoid excessive loading of shoulder
  - No heavy lifting
  - No prolonged, repetitive upper limb activities

#### Milestones at 14 Weeks:

- Resolved pain, inflammation and muscle inhibition
- AROM with good SHR through elevation to 100% of pre-operative range
- Passive external rotation to 100% of pre-operative range

#### Phase 4 (3 – 6 Months)

##### Goals:

- Maintain full PROM
  - Continue multi-directional stretching into end of range **as tolerated/not into pain**
  - Capsular stretches (especially posterior capsule if tight)
- Full pre-operative AROM with good SHR **as tolerated/not into pain**
- Progress shoulder strengthening **as tolerated/not into pain**
  - Progress cuff strengthening and scapular stabilisation exercises
    - Progress proprioceptive exercises
    - Progress shoulder and upper limb strengthening ensuring good SHR
    - Progress dynamic and rhythmic stabilisation exercises
- Progress biceps strengthening
- Progress lower limb and core strengthening as required
- Gradually progress functional activities
- Gradually progress sport-specific exercises

##### Precautions:

- Avoid excessive loading of shoulder
  - No heavy lifting away from body
  - No prolonged, repetitive overhead activities

### Milestones at 6 months:

- Full pain-free motion and rotator cuff strength restored

### Phase 5 (6 – 9 Months)

#### Goals:

- As phase 4
- Gradual return to strenuous work activities as required
- Gradual return to contact sports as required

---

To contact the Consultant Secretary, call 07810356433 or to contact the Physiotherapy team for Sulis Hospital call 01761 422388, for Physiotherapy Circle Health Group Bath Clinic call 01225 838767, for Physiotherapy St Joseph 01633 820321